

WILL THE REAL HYALURONAN PLEASE STAND UP?

Luke R. Bucci, Ph.D., CNS, CCN and Amy A. Turpin, M.Sc.

This review intends to inform readers about the identity of Hyaluronan (Hyaluronic Acid or HA) materials currently used in dietary supplement products. As contestants from the TV game show “To Tell The Truth” (1956-1978) were made to choose the real person from two other foils by asking questions of each, this review will answer questions so that the real HA can stand up. There are major differences between the three major types of HA materials used in dietary supplements. According to the voluminous scientific literature, these differences may lead to distinctly different biological effects. This review does not attempt to discuss the rationale for oral HA as opposed to injectable or topical uses – this is an obvious topic for further review. However, ingestible HA products are available to the general public, and their use is growing to the point that HA-containing products are available from almost every mass-market retail provider. Consumers and health care professionals alike are not given facts about the identity of HA materials used in dietary supplements. Informed choices for evaluation of HA-containing products cannot be made without an understanding of what materials are used for each product.

HA PRIMER

HA is an extremely large molecule (normally 1-5 million daltons) of repeating subunits of N-acetylglucosamine and glucuronate (1-14). HA is classified as a glycosaminoglycan (GAG), and is present physiologically and commercially as the sodium salt (sodium hyaluronate). Single HA molecules exist as a hydrophilic, coiled mass approximately 200 nm in diameter, but form a mesh under typical *in vivo* conditions (8). HA is found everywhere in the body, but has especially important roles in joints. HA is the primary agent responsible for the viscosity and lubricating properties of synovial fluid (6, 8, 11-29). In cartilage, HA serves as the backbone of aggrecan proteoglycans, bound to core and link proteins (18, 30-49). Furthermore, HA is an impor-

tant and under-recognized regulator of cell and tissue functions, as shown by the presence of specific receptors on all cells for HA and its fragments (8, 11-14, 50-65).

The scientific and medical literature on HA and its effects on joints has had virtually no references to oral use of HA until very recently. Instead, HA has a prodigious literature on biological effects (6, 8, 11-14, 19-26, 28, 62, 66, 67), including a rather large number of *in vivo* animal and human clinical studies using intraarticular injections to treat osteoarthritis (OA) with purified HA (reviewed in 21, 22, 25, 26, 68-86). Thus, there is substantial knowledge of the biological effects of purified, high molecular weight HA on joint health.

HYALURONAN AS AN ORAL DIETARY SUPPLEMENT

HA as an oral supplement entered mainstream consciousness after an ABC News pro-

gram aired a segment that attributed possible anti-aging effects to dietary HA in November 2002. Since then, HA has been appearing as an ingredient in an ever-increasing number of dietary supplement products. HA-containing products have focused on joint and skin health. At the beginning of 2005, several of the largest dietary supplement companies were selling products containing HA in major retail outlets throughout the United States. This is in addition to other products available through Internet sales. In short, HA has catapulted from obscurity to mainstream store shelves rapidly.

There are a few animal studies on joint health after intravenous or intraperitoneal administration of HA (87-89), which provide some insight into tissue uptake and possible effects of HA after its appearance in the bloodstream. Four recent reports investigated effects of oral HA on animal musculoskeletal tissues. One report from the Hyaluronan 2003 Proceedings by Matrix Biology Institute administered 100 mg HA orally to 12 racing thoroughbreds for 59 days (90). Horses given HA were examined for lameness less frequently than 13 control group horses, suggesting that oral HA prevented lameness in active horses. Another report by Stancikova *et al* administered HA (0.5-1 mg/kg) with molecular weights of 0.75 and 1.62 million daltons by oral routes for eight weeks to ovariectomized rats (91). Both sizes of HA showed increased serum concentrations of nitric oxide (NO), but only the larger HA showed decreased markers of bone resorption as well as an inhibition of bone mineral density loss. Two other animal studies were found on the Contipro website (92). Oral HA fed to rats at a dose of 0.1 mg/kg showed reduced swelling of hind leg and decreased NO production after arthritis induction by Freund's adjuvant. An open study from 7 veterinary clinics studied oral HA given to 53 racehorses with inflammatory and degenerative joint conditions. After 30 days of oral HA, symptoms were reduced and functionality improved. All reports used purified, high-mole-

cular weight HA from a microbial fermentation source (Nutrihyl®, Contipro Group Holding, Czech Republic).

There are unpublished reports of oral HA in humans for joint health from the Czech Republic. A product named Chondrorevit was reported to improve joint health in subjects with knee OA or after knee surgeries (92). Daily doses of 20 mg HA and 400 mg chondroitin sulfate were administered for 90 days. Improvements in joint function or recovery time after surgery were reported. Interestingly, this amount of chondroitin sulfate has not been shown to have significant efficacy in dose-finding studies from Europe (93), and thus, any benefits may be attributable to the HA or the combination of HA and chondroitin at a less than efficacious dose. However, these studies did not appear to be randomized, double blind or placebo controlled, and thus, their findings must be verified in controlled studies before results can be relied upon. Nevertheless, HA has been reported to have benefits for joint health after oral use in animal and human clinical studies. At this point in time, reports of oral HA for joint health have all used high molecular weight, purified HA, except for one study discussed later in this review.

MOLECULAR WEIGHT AND BIOLOGICAL EFFECTS OF HYALURONAN: EVIDENCE SHOWING THAT SMALL HYALURONAN AND FRAGMENTS ARE DIFFERENT

Because HA is a repeating polymer devoid of modifications (such as branching or sulfation), major differences in biological activities for HA are attributable to its molecular weight (size) (25, 65, 67, 94-98). There are clear cutoffs of molecular weight for lubricating properties (17, 19, 25, 99-106), cell signaling (107-117), and perhaps for efficacy of injectable intraarticular preparations for OA treatment (63, 67, 118-120). Molecular weight of less than 500,000 daltons seems to be

the cutoff for differing properties of HA. This concept is summarized by a comment from Camenisch and McDonald: “The work of Ohkawara and coworkers and numerous other observations suggest that there is something fundamentally different about the biological response to high (megadaltons) and lower molecular weight HA.” (95). In essence, bigger is better for health benefits from HA, and smaller fragments have very different or opposite properties than native, high molecular weight HA.

Very low molecular weight HA fragments cannot be expected to have the same biological effects as high molecular weight HA, as is clear from the growing literature on biological effects of HA. Very low molecular weight fragments either do not bind to specific cell membrane receptors that recognize native HA, or they transduce different signals to cells (65, 107, 108, 113, 121-126). For example, very low molecular

weight HA fragments (6.9 kDa), but not high molecular weight HA, have been associated with promotion of cancer cell neovascularization, migration and metastasis (112, 117, 122, 127-132). Low molecular weight HA fragments have either lost anti-inflammatory effects or instigate proinflammatory, catabolic properties compared to native HA in several biological systems (64, 110, 114, 116, 121, 125, 132-141). For example, HA fragments of 250,000-dalton molecular weight induced expression of inflammatory genes in macrophages (64, 65, 132, 135, 142). These effects are not due to contamination, as shown by the fact that the size range of HA tested *in vitro* is the same range as is found *in vivo* during inflammation (143). Some reviewers have postulated that HA fragments promote and propagate OA (65, 97, 136, 140, 144). These are obviously not desirable traits for dietary supplements purporting to support joint health.

TABLE 1: Identity and Characteristics of Hyaluronan Materials Used in Dietary Supplements

Hyaluronan Type	Source	Molecular Weight (daltons)	Purity	References
Hydrolyzed chicken sternal cartilage Type II Collagen	Chicken sternal cartilage	50–10,000 ¹	~10%	<ul style="list-style-type: none"> • Ishaq S. Hyaluronic acid and chondroitin sulfate based hydrolyzed collagen type II and method of making same. United States Patent 6,780,841, August 24, 2004. • http://www.biocelltechnology.com
Rooster Comb (Injuv™)	Chicken comb	50,000–200,000	9%	<ul style="list-style-type: none"> • Udell RG, Naguib YMA. Hyaluronic acid in soft gel form. United States Patent 6,806,259, October 19, 2004. • http://www.injuv.us
Sodium hyaluronate	Microbial fermentation (Streptococcus species) or chicken comb	700,000–1 million or higher	90+%	<ul style="list-style-type: none"> • http://www.bioiberica.com/jointcare/hyaljoint.htm • http://www.cpn-contipro.com

¹Patent 6,780,841 (column 6, line 12) stated: “The average molecular weight of the final product is between 50 and 10,000 daltons, preferably 5,500 daltons.” Neither the patents nor website specifically list a range of molecular weight for hyaluronan itself. Methods of analysis to determine molecular weight or identity of HA were not stated.

HA MATERIALS

There are three major types of commercially available HA materials used in dietary supplements. Table 1 lists these types and their basic attributes. Information was obtained in the public domain from material supplier or supplement company websites. It is readily apparent that there are major differences in molecular weight of each type, yet all are labeled identically as Hyaluronic Acid or Sodium Hyaluronate on dietary supplement product labels (see Table 2). Only one type of HA matches what is normally present in the human body – high molecular weight, purified (protein-free) HA. The other two major types of HA materials have lower molecular weights.

HYDROLYZED CARTILAGE HA (BIOCELL COLLAGEN II™)

HA from hydrolyzed cartilage has a patent describing the enzymatic degradation of Type II collagen from chicken sternal cartilage into very small fragments (50 to 10,000 daltons), with no high molecular weight material remaining (145). This process results in a material consisting of 10% Hyaluronic Acid, 20% depolymerized chondroitin sulfate, and around 63% Type II collagen fragments (145). This review is not concerned with the collagen or chondroitin contents of this material, which may have their own effects on joint health. However, the HA in this material is pertinent since it is used in dietary supplement products listed as a source of HA (see Table 2). It is not stated in the patent or in company website information what size range of HA fragments exist in this material. The patent further states (column 7, line 46):

“By extracting HA from chicken sternal cartilage, a low molecular weight HA may be obtained by hydrolysis after such extraction. This distinction is crucial because the beneficial therapeutic activity of HA is mostly

dependent upon the molecular weight of HA. Due to the low molecular weight of the HA found in the hydrolyzed collagen type II, the hydrolyzed collagen type II readily absorbs into the gastrointestinal tract and allows the rejuvenating constituents of HA and CSA to restore viscoelasticity to the skin, protect connective tissues, promote cartilage synthesis, retain skin moisture, heal wounds and improve the overall appearance of skin.”

The patent does not supply any data on the size of HA, the analytical method used to assess HA identity, or any effects of the material. In other words, there is no supportive data to substantiate the properties attributed to HA in the statement above. However, it is clear that the inventor assumes that the HA in this material is of low molecular weight.

As briefly discussed previously in this review, HA fragments do not have the same properties in biological systems as native, high-molecular weight HA does, and thus, this material does not fit the definitions and descriptions of HA agreed upon by textbooks, reviewers and experts in the field (8, 11-14, 19, 23-26, 28, 62, 113, 125, 126). In short, this material is not typical HA and thus should not be described as HA on dietary supplement labels. Rather, this material is more accurately described as very low molecular weight HA oligosaccharides or even pre-digested HA. This statement assumes that the HA in this material is indeed hydrolyzed, which at present is not substantiated.

Furthermore, HA in cartilage does not exist as free, unbound molecules, but rather, as the backbone of aggrecan proteoglycans (18, 30-49). As such, HA is bound to link and core proteins, and sometimes to other GAGs or chondrocytes themselves. Even assuming proteolytic enzyme treatment as described in the patents to prepare this material can cleave HA from protein (145-147), the presence of amino acids crosslinked to HA fragments are not synonymous with a definition of HA.

TABLE 2. Examples of How Dietary Supplement Products Containing Hyaluronan Are Labeled¹

Product	Description of Hyaluronan in Supplement Facts	Amount per serving	Accompanying Label Claims
21 st Century Healthcare, Inc. Arthri-Flex	Hyaluronic Acid	Not stated—part of a Flexicol TM) proprietary blend with four other ingredients totaling 645 mg	<ul style="list-style-type: none"> • With Flexicol Premium Chicken Sternum Type II Collagen • Containing: Premium Chicken Sternum Cartilage Type II Collagen (naturally occurring Chondroitin Sulfate and Hyaluronic Acid) • Contains Highly Absorbable Hyaluronic Acid • www.21stcenturyvitamins.com
Arthritis Research Corp. Flex-a-min [®] Complete TM	Hyaluronic Acid (as Sodium Hyaluronate)	Not stated—part of a proprietary blend with three other ingredients totaling 400 mg	<ul style="list-style-type: none"> • Hyaluronic Acid—The Glucosamine in Flex-a-min[®] CompleteTM promotes the production of Hyaluronic Acid in the body, including the synovial fluid surrounding joints. • www.flexamin.com
Arthritis Research Corp. Flex-a-min [®] Triple Strength	Hyaluronic Acid (as Sodium Hyaluronate)	Not stated—part of proprietary blend with Silica totaling 40 mg	<ul style="list-style-type: none"> • Plus, we added the popular ingredient Hyaluronic Acid to set this formula apart. • Hyaluronic Acid—The Glucosamine in Flex-a-min[®] CompleteTM promotes the production of Hyaluronic Acid in the body, including the synovial fluid surrounding joints. • www.flexamin.com
Injuv TM Hyaluronic Acid	Hyaluronic Acid	12.6 mg (from 140 mg Injuv TM) ²	<ul style="list-style-type: none"> • Stop the stiffness, stop the pain and increase your mobility with InjuvTM (www.jointlubricant.com/injuvworks.htm) • The HA in InjuvTM stops pain and inflammation of joints, because it helps to hydrate and lubricate the synovial cell membrane. (www.jointlubricant.com/injuvworks.htm) • InjuvTM has an exclusive patent on the process that breaks down HA so that it can be taken orally. • The hyaluronic acid molecule in Injuv has been enzymatically cleaved into smaller fragments. (www.injuv.us)
Nature's Bounty Hyaluronic Acid	Hyaluronic Acid (as Sodium Hyaluronate)	20 mg per capsule	<ul style="list-style-type: none"> • Hyaluronic Acid is found in almost all adult connective tissue. Hyaluronic acid is a natural component of the body. • www.NaturesBounty.com
Schiff [®] Move Free Triple Strength Product	Hyaluronic Acid (in Other ingredients list)	Not stated (Consumer Service line stated 3.3 mg per serving)	<ul style="list-style-type: none"> • Now with Joint Fluid (Hyaluronic Acid) • www.schiffvitamins.com

TABLE 2. Continued

Product	Description of Hyaluronan in Supplement Facts	Amount per serving	Accompanying Label Claims
Spring Valley Glucosamine & Collagen with Chondroitin & HA (Hyaluronic Acid)	Hyaluronic Acid	40 mg	<ul style="list-style-type: none"> • In a base of amino acids (Hydrolyzed Collagen Type II) • Collagen Type II 500 mg • Chondroitin Sulfate (Avian, Porcine, Bovine) 100 mg • Collagen is the most plentiful protein and building block in the body and is a natural source of Chondroitin, a nutrient shown to regenerate joint tissue, and Hyaluronic Acid (HA), a revolutionary compound that retains moisture and functions as a lubricant between connective tissues.* • Hyaluronic acid is fast absorbing in the blood and thus available for joint tissues. • Lot number 4MA0236
Swanson Ultra® Hyal-Joint™	Hyal-Joint™ Hyaluronic Acid	20 or 50 mg per capsule.	<ul style="list-style-type: none"> • The unique hyaluronic acid formulation in Hyal-Joint has the proven ability to be absorbed in the intestinal tract and has demonstrated significant joint-protective properties in numerous studies. • www.bioiberica.com
Windmill™ Health Products Glucoflex™ Hyaluronic Acid	Hyaluronic acid sodium salt	10 mg per tablet	<ul style="list-style-type: none"> • Advanced Joint Lubricating Formula with Glucosamine • Glucoflex™ Hyaluronic Acid formula with Glucosamine supplies the most bioavailable form of HA (Hyaluronic Acid) for the most direct delivery of this amazing nutrient.

¹Products were obtained from store shelves from the United States or from internet website information. Products are listed in alphabetical order according to manufacturer name.

²Some products containing Injuv™ HA list the total amount of material instead of the actual amount of HA, misleading consumers into believing there is 70 or 140 mg of HA per softgel.

Another issue apparently not considered by some marketers of hydrolyzed cartilage HA is the integrity of borrowing literature describing properties of HA to support such product claims (see Table 2). Since hydrolyzed HA has quite different physical and biological properties compared to high molecular weight HA, it is a different entity. For products with very low molecular weight HA (hydrolyzed cartilage) it is inappropriate to make claims, such as lubricating properties, which rely on literature describ-

ing high molecular weight HA. This is especially true when there is also evidence of very low molecular weight HA not having the properties claimed (such as lubricating properties). Thus, it would be more appropriate to base any claims for joint health with this material on studies using this material.

Recently, a late-breaking abstract presented at the 2004 FASEB meeting in Washington DC described the results of a human clinical study using hydrolyzed cartilage material (148). A press

release from BioCell Technology, LLC posted on their website on April 2003 also reiterated the results of the abstract (149). In the abstract, the material was described as “. . . a hydrolyzed type II collagen product (avian source: Biocell™; HC2)” without mention of HA. A dose of 2000 mg daily was given to eight subjects with knee or hand osteoarthritis for two months, and compared to a placebo group of eight subjects in a randomized, double blind, placebo-controlled study. WOMAC scores were improved significantly greater in the HC2 group compared to placebo, but no data were presented in the abstract or press release. This amount of material would contain approximately 200 mg of hydrolyzed HA daily, according to the supplier's website and patent. However, approximately 1600 mg of hydrolyzed Type II collagen and 400 mg of chondroitin sulfate fragments were also present in this material. Thus, the contribution of HA to observed results is unclear. Furthermore, human studies with subjective questionnaire endpoints (WOMAC) and with only eight subjects per group run a high risk of low statistical power, making results preliminary until confirmed in more powerful studies. Most other studies that mixed subjects with knee and hand osteoarthritis have been criticized since the arthritic process in hand osteoarthritis is distinctly different than in knees.

In addition, a press release by BioCell Technology from February 2, 2004 described a 36-hour peak absorption study conducted in an unknown number of human subjects with hydrolyzed cartilage material (150). HA levels were determined after a single dose and after 28 days of administration. Levels of HA in blood were claimed to be 7008.62% above control levels after 12 hours. Also, two HA metabolites 1/600th the size of the ingested HA were found. After 28 days, HA levels in blood were reported to be 3542.58% above control, and metabolites at 11,890.15% above control. No details on blood collection or analytical methods were given, although the reporting of HA metabo-

lites with two decimal places implies a high degree of precision for the method of analysis. Normal human serum levels of HA are 0.01-0.1 mg/L (mcg/ml) (8). Assuming the press release was describing serum HA levels, a 7000% increase in blood HA levels would produce levels of 0.7-7 mcg/ml, which is within the range seen for certain cancers, liver conditions and inflammatory diseases (8). One possible interpretation of this study is that this material elevates serum levels of HA fragments, along with native HA levels, although more experimental details are needed to confirm this interpretation because the effects of activity and food intake are known to increase plasma HA levels (151). Absorption of HA from the material is not proven by these results, since HA can rapidly enter circulation, especially in inflammatory diseases (8). Thus, another interpretation of these results is that this material is pro-inflammatory, which in and of itself would elevate serum levels of HA, congruent with findings of other researchers on pro-inflammatory effects of very low molecular weight HA fragments.

Assay of HA fragments, especially in a mixture with chondroitin sulfate or its hydrolysis products, is problematic. In fact, assay of products containing very low molecular weight HA fragments by validated assays utilizing HA binding protein technology do not find any measurable HA (Bucci, L., unpublished results). This indicates that if present, HA fragments are less than eight subunits in length. These assays also indicate that there is no high molecular weight HA present, so claims for this material on product labels or websites based on properties of high molecular weight HA cannot be substantiated by borrowing from the literature describing high molecular weight HA.

Therefore, HA derived from hydrolyzed chicken sternal cartilage is characterized by extremely low molecular weights, a lack of identity as native HA itself, a probable difference in biological effects (assuming uptake and distribution to tissues), a lack of transparency on how

the HA content was verified, and inability to show presence of HA in validated assays for HA using specific HA binding protein methodology. In addition, some products do not describe this material as hydrolyzed (see Table 2), which misleads consumers into thinking that they are getting native HA. Consumers expect to receive high molecular weight HA since this is what media, medical usage and website information all describe as characteristic of HA. This material is easy to identify in products because of the presence of Type II Collagen, chicken (avian) sternal cartilage source, and simultaneous presence of HA, chondroitin sulfate and Type II collagen naturally. These facts argue strongly that this material should not be labeled as HA in dietary supplement products as the identity and putative biological effects are much different from that of high molecular weight HA.

ROOSTER COMB HA (INJUV™)

The second major type of HA used in dietary supplement products is prepared from dried rooster comb, a rich natural source of HA (7500 mg/L) (8). This material is claimed to be enzymatically denatured, with a molecular weight of 50,000 to 200,000 daltons in order to facilitate uptake, without any supporting evidence (152). Injuv™ HA is identified by validated HA binding protein assays (Bucci L, unpublished data). However, experts in the field regard HA with a molecular weight of 50,000-200,000 daltons to be low molecular weight HA. Thus, HA in this size range has lost its ability to form networks in solution, and thus, has lost viscosity and lubricating properties.

Inspection of HA in intact rooster combs shows tight association of HA with proteins (153-155). It is unclear if conditions used to prepare rooster comb HA with enzymatic hydrolysis would actually dissociate the tightly bound proteins from the HA, since details of preparation are lacking in the published patent. High molecular weight HA prepared from rooster combs for

injectable use, such as Hyalgan® Sodium Hyaluronate from Sanofi-Synthelabo Inc., has been extensively treated to remove proteins in order to prevent reactions after injection, and has not shown appreciable adverse effects after intraarticular injections (156). Such material would fit into the third type of HA material.

HA with molecular weights of less than 500,000 daltons exhibit decreased viscosity and lubricating properties in solution, as stated earlier in this review. HA in this size range has lost its ability to form networks in solution. Therefore, claims about lubricating properties are inappropriate with this material unless specific studies are performed to investigate its lubricating effects in joints. Also, signals transduced by HA in the molecular weight range of 100,000-200,000 are different from those of HA at 1 million daltons, and generally are proinflammatory. The presence of tightly bound proteins in rooster comb HA has uncertain implications for uptake of this type of HA. Digestion and uptake of rooster comb HA may be different from native, purified HA. So this material, while clearly low molecular weight HA, is different from native, high molecular weight HA and has been shown to have different biological effects.

PURIFIED HA

The third major type of HA available for dietary supplement products is essentially purified HA (Table 1). Typical commercial sources are from microbial fermentation (*Streptococcus* species) or rooster combs. Molecular weight is around 1 million daltons, which corresponds to native HA in cartilage and in synovial fluid. Protein is absent or very low in these preparations. HA from microbial fermentation sources can be declared as vegetarian, whereas HA derived from any cartilage or chicken comb sources cannot. Since purified HA for food use is produced by manufacturers that also produce purified HA for pharmaceutical use, there is more certainty that these products have a tradition of quality

TABLE 3. Biological Properties of High Molecular Weight Hyaluronan and Hyaluronan Fragments

Molecular Weight (daltons)	Biological Property	Reference
Very low molecular weight oligosaccharides (800-5000)	<ul style="list-style-type: none"> • Angiogenic—Induced angiogenesis in rat skin grafts, healing wounds, chick chorioallantoic membranes • Angiogenic—Induced rapid transient up-regulation of the immediate early genes c-fos, c-jun, jun-B, Krox-20 and Krox-24 that control angiogenesis • Angiogenic—Increased tyrosine kinase, protein kinase C cascades in endothelial cells • Angiogenic—Induced proliferation of bovine brain and aortic endothelial cells • Antiinflammatory—Tetrasaccharides, but not larger sizes of HA, induced heat shock protein 72 in K562 cells and prevented cell death of PC12 cells after serum deprivation. • Antioxidant—Exhibited antioxidant activity against superoxide, but less than higher molecular weight HA; did not scavenge hydroxyl radicals (unlike high molecular weight HA) • Catabolic—Did not affect sulfate incorporation in chondrocyte cultures • Catabolic—Blocked binding of aggrecan HA to chondrocytes, resulting in loss of staining for proteoglycans (disrupted cartilage structure) • Catabolic—Caused chondrolysis (loss of chondrocytes) when hexasaccharides exposed to cartilage tissue slices and chondrocyte cultures • Inflammatory—High molecular weight HA cleaved at sites of inflammation into small fragments • Inflammatory—Activated macrophages and dendritic cells in healing wounds • Inflammatory—Induced maturation of dendritic cells to tissue macrophages • Inflammatory—Induced production of IL-1beta, TNF-alpha, IL-12 and iNOS in dendritic cells and macrophages • Inflammatory—Did not inhibit arachidonic acid release after bradykinin from synovial fibroblasts from osteoarthritic subjects • Inflammatory—Did not inhibit MMP-1 and RANTES expression or production in normal and osteoarthritic human chondrocytes in culture • Inflammatory—Induced expression of matrix metalloproteinases, but not TIMPs, in normal rabbit and bovine cartilage chondrocytes • Inflammatory—Induced MMP-9, MMP-13 expression and production in murine fibroblasts and 3LL cells • Inflammatory—Stimulated MAP kinase and urokinase-type plasminogen activator in human HCS-2/8 chondrosarcoma cells • Inflammatory—Did not inhibit neutrophil adhesion or aggregation under conditions of shear and turbulent flow • Inflammatory—Stimulated tyrosine phosphorylation and c-Met expression in human chondrosarcoma cells 	7, 8, 11, 15, 20, 28-32, 35, 38, 39, 42, 46, 52, 53, 57, 59, 60-63, 67-69, 71-73

TABLE 3. Biological Properties of High Molecular Weight Hyaluronan and Hyaluronan Fragments (*continued*)

Molecular Weight (daltons)	Biological Property	Reference
24,000	<ul style="list-style-type: none"> • Catabolic—Inhibited tumor growth, induced apoptosis • Inflammatory—Induced expression and production of inflammatory mediators from murine macrophages (IL-1, TNF-alpha, IL-12, iNOS, metalloelastase) • Lubrication—Reduced elasticity & viscosity of 1.3MDa HA solutions (competed for meshwork formation) 	13, 23, 36, 37, 43, 44, 54
33,000	<ul style="list-style-type: none"> • Antiangiogenic—Did not induce angiogenesis of rat skin grafts 	35
<50,000	<ul style="list-style-type: none"> • Catabolic—Did not stimulate HA synthesis or inhibited synthesis (high concentrations) of HA from human osteoarthritic synovial fibroblasts • Inflammatory—Did not inhibit proliferation of human mononuclear cells 	47, 58
50,000	<ul style="list-style-type: none"> • Inflammatory—Did not inhibit MMP-1 and RANTES expression or production in normal and osteoarthritic human chondrocytes in culture • Inflammatory —Induced expression and production of inflammatory mediators from murine macrophages (IL-1, TNF-alpha, IL-12, iNOS, metalloelastase) 	23, 36, 37, 43, 44, 61
90,000	<ul style="list-style-type: none"> • Catabolic—did not enhance bone mineralization in rat calvarial cells • Inflammatory—Did not inhibit phagocytosis by mouse peritoneal macrophages at same viscosity as higher molecular weight HA • Lubrication—Osmotic pressure opposing fluid drainage in rabbit joints reduced, facilitates escape of fluid and HA from joints 	6, 10, 24
80,000–200,000	<ul style="list-style-type: none"> • Antiinflammatory—Did not induce matrix metalloproteinase production in murine fibroblasts and 3LL cells • Antiinflammatory—Did not induce IL-1beta, TNF-alpha or IL-12 in human dendritic cells or macrophages • Antioxidant—Scavenged hydroxyl radicals in vitro, but was less effective than high molecular weight HA for protecting tendon fibroblasts from hydroxyl radicals • Inflammatory—Did not inhibit cell proliferation of rabbit synovial cells • Inflammatory—Induced expression and production of inflammatory mediators from murine macrophages (IL-1, TNF-alpha, IL-12, iNOS, metalloelastase) 	8, 16, 23, 36, 37, 43, 44, 49, 62, 63
200,000	<ul style="list-style-type: none"> • Inflammatory—Markedly stimulated iNOS mRNA in liver endothelial and Kuppfer cell types • Inflammatory—Intravenous HA (250,000 daltons) did not inhibit proinflammatory cytokine levels in serum after liver injury in rats • Inflammatory—In human eosinophils, increased production of intercellular adhesion molecule-1, TGF-beta, protein secretion, and transformed cells from round to spindle shapes 	19, 40, 45, 51

TABLE 3. Biological Properties of High Molecular Weight Hyaluronan and Hyaluronan Fragments (*continued*)

Molecular Weight (daltons)	Biological Property	Reference
	<ul style="list-style-type: none"> • Inflammatory—Human peritoneal mesothelial cells produced more inflammatory cytokines (MCP-1, IL-8) 	
280,000	<ul style="list-style-type: none"> • Anticatabolic—Inhibited cartilage degradation by neutrophils, but significantly less than 950,000 or 2.0 million dalton HA • Inflammatory—Produced inflammatory cytokines RANTES, IL-12, MIP from murine macrophages 	21, 65
350,000	<ul style="list-style-type: none"> • Lubrication—Formed poor meshworks with reduced viscosity • Lubrication—Osmotic pressure opposing fluid drainage in rabbit joints reduced, facilitates escape of fluid and HA from joints 	6, 54
460,000–2.8 million	<ul style="list-style-type: none"> • Antiinflammatory—Dose-dependent inhibition in phagocytosis from mouse peritoneal macrophages (steric hindrance) • Antiinflammatory—Did not induce matrix metalloproteinases in normal rabbit and bovine cartilage chondrocytes • Antinociceptive—Molecular weight-dependent improvement in abnormal gait in rats acute arthritis, indicating nociceptive effects with higher molecular weight HA 	1, 10, 17, 46
Native HA (approx. 800,000 to 6 million)	<ul style="list-style-type: none"> • Anabolic—Stimulated synthesis of high molecular weight HA from human osteoarthritic synovial fibroblasts • Anabolic—Increased proteoglycan synthesis from IL-1Beta treated chondrocytes • Anabolic—Stimulated fibroblast proliferation in a collagen lattice, but not in fibroblast monolayer • Anabolic—Enhanced proteoglycan synthesis in human cartilage explants after fibronectin fragment treatment • Anabolic—Enhanced chondrocyte proliferation and chondroitin sulfate synthesis in rabbit chondrocytes embedded in collagen gels • Anabolic—Forms pericellular matrix (with aggrecan) in presence of chondrocytes • Anabolic—Improved bone mineralization in rat calvarial cells • Antiangiogenic—Inhibited stimulation of angiogenesis and early response genes by low molecular weight HA fragments • Antiangiogenic—Inhibited or did not induce angiogenesis of chick chorioallantoic membranes or endothelial cells • Antiangiogenic—Did not cause proliferation of bovine brain and aortic endothelial cells • Anticatabolic—Decreased reduction in proteoglycan synthesis from fibronectin fragments • Anticatabolic—Decreased cartilage degradation from fibronectin fragments • Anticatabolic—Decreased expression of matrix metalloproteinase 3 in human cartilage explants after fibronectin fragment treatment • Anticatabolic—Inhibited leukocyte elastase and lysozyme release • Anticatabolic—Inhibited neutrophil-mediated cartilage degradation (reduced release of sulfated GAGs from bovine nasal cartilage explants) • Antiinflammatory—Did not induce iNOS mRNA from liver cell types 	1-8, 10, 12, 15-18, 20-22, 24-29, 32-34, 38, 39-44, 45, 47-51, 53, 58, 59-63, 66, 68, 70-72, 74, 75

TABLE 3. Biological Properties of High Molecular Weight Hyaluronan and Hyaluronan Fragments (*continued*)

Molecular Weight (daltons)	Biological Property	Reference
	<ul style="list-style-type: none"> • Antiinflammatory—Reduced IL-1beta, TNF levels in rat air pouches • Antiinflammatory—Did not induce IL-1beta, TNF-alpha or IL-12 in human dendritic cells or macrophages • Antiinflammatory—Inhibited PGE2 production by human osteoarthritic synovial cells and normal rabbit articular chondrocytes after IL-1alpha induction • Antiinflammatory—Did not induce differentiation of human dendritic cells into macrophages • Antiinflammatory—Decreased mitogen-dependent activation of human mononuclear lymphocytes • Antiinflammatory—Inhibited proliferation of rabbit synovial cells in culture • Antiinflammatory—Inhibited carrageenin-induced edema in dose-dependent manner • Antiinflammatory—Inhibited adjuvant arthritis in dose-dependent manner • Antiinflammatory—Inhibited arachidonic acid release induced by bradykinin in synovial fibroblasts of osteoarthritic humans • Antiinflammatory—Inhibited advanced glycation endproducts from activating NF-kappaB, IL-1alpha, IL-6, TNF-alpha • Antiinflammatory—Inhibited MMP-1, RANTES expression and production in normal and osteoarthritic human chondrocyte cultures • Antiinflammatory—Did not enhance MAP kinase, urokinase-type plasminogen activator, tyrosine phosphorylation or c-Met in human HSC-2/8 chondrosarcoma cells • Antiinflammatory—Enhanced production of TIMP-1 in normal bovine chondrocyte culture, lowered stromelysin/TIMP-1 ratios • Antiinflammatory—Decreased synovial PGE2 and bradykinin in arthritic rats after intraarticular injection • Antiinflammatory—Inhibited neutrophil aggregation and adhesion under conditions of shear and turbulent flow • Antiinflammatory—Intravenous HA (>780,000 daltons) decreased proinflammatory cytokines and liver enzymes after liver damage in rats • Antiinflammatory—Did not cause eosinophils to produce inflammatory cytokines • Antiinflammatory—Did not induce inflammatory cytokines RANTES, IL-12, MIP from murine macrophages • Antioxidant—Decreased free radical formation from IL-1Beta treated chondrocytes • Antioxidant—Scavenged superoxide and hydroxyl radicals dose-dependently • Antioxidant—Protected tendon fibroblasts in culture from damage by hydroxyl radicals • Lubrication—High molecular weight HA (>500,000 daltons) caused outflow buffering in rabbit joints, reducing drainage and loss of HA and fluid • Regulatory—Inhibited sulfate incorporation into chondrocyte cultures 	

TABLE 3. *Continued*

Note: Filion and Phillips determined that some (2/7) of both high (0.5-2 million daltons) and low (10,000-300,000) molecular weight preparations of HA were inflammatory (produced IL-12 and TNF from human monocytes) due to contaminating DNA (based on reduction of activity after DNase treatment). This suggested that a low percentage of individual HA preparations have potential to cause inflammatory events not due to the molecular weight of HA. However, many of the studies cited in Table 3 examined different molecular weights from the same source of HA, and noticed an effect dependent on molecular weight, indicating contamination was not involved. The large number of studies showing an effect dependent on molecular weight from a wide variety of investigators, preparations and biological sources also argues strongly that DNA contamination is not operative in at least a majority of studies, or else no differences dependent on molecular weight would have been found. Scott and Heatley provide tertiary structure evidence that smaller HA fragments are different from high molecular weight HA, helping to explain the differences in biological activities of denatured HA.

References for Table 3:

- 3.1 Aihara S, Muarakami N, Ishii R, Kariya K, Azuma Y, Hamada K, Umemoto J, Maeda S. [Effects of sodium hyaluronate on the nociceptive response of rats with experimentally induced arthritis.] *Nippon Yakurigaku Zasshi* 1992 Oct; 100(4):359-365.
- 3.2 Akatsuka M, Yamamoto Y, Tobetto K, Yasui T, Ando T. In vitro effects of hyaluronan on prostaglandin E2 induction by interleukin-1 in rabbit articular chondrocytes. *Agents Actions* 1993 Jan; 38(1-2):122-125.
- 3.3 Akatsuka M, Yamamoto Y, Tobetto K, Yasui T, Ando T. Suppressive effects of hyaluronic acid on elastase release from rat peritoneal leukocytes. *J Pharm Pharmacol* 1993 Feb; 45(2):110-114.
- 3.4 Anastassiades T, Robertson W. Modulation of mitogen-dependent lymphocyte stimulation by hyaluronic acid. *J Rheumatol* 1984 Dec; 11(6):729-734.
- 3.5 Camenisch TD, McDonald JA. Hyaluronan: is bigger better? *Am J Respir Cell Mol Biol* 2000 Oct; 23(4):431-433.
- 3.6 Coleman PJ, Scott D, Mason RM, Levick JR. Role of hyaluronan chain length in buffering interstitial flow across synovium in rabbits. *J Physiol* 2000 Jul 15; 526 (Pt 2):425-434.
- 3.7 Deed R, Rooney P, Kumar P, Norton JD, Smith J, Freemont AJ, Kumar S. Early-response signalling is induced by angiogenic oligosaccharides of hyaluronan in endothelial cells. Inhibition by non-angiogenic, high-molecular-weight hyaluronan. *Int J Cancer* 1997 Apr 10; 71(2):251-256.
- 3.8 Fieber C, Baumann P, Vallon R, Termeer C, Simon JC, Hofmann M, Angel P, Herrlich P, Sleeman JP. Hyaluronan-oligosaccharide-induced transcription of metalloproteases. *J Cell Sci* 2004 Jan 15; 117(Pt 2):359-367.
- 3.9 Filion MC, Phillips NC. Pro-inflammatory activity of contaminating DNA in hyaluronic acid preparations. *J Pharm Pharmacol* 2001 Apr; 53(4):555-561.
- 3.10 Forrester JV, Balazs EA. Inhibition of phagocytosis by high molecular weight hyaluronate. *Immunology* 1980 Jul; 40(3):435-436.
- 3.11 Forrester JV, Lackie JM. Effect of hyaluronic acid on neutrophil adhesion. *J Cell Sci* 1981 Aug; 50:329-344.
- 3.12 Fukuda K, Takayama M, Ueno M, Oh M, Asada S, Kumano F, Tanaka S. Hyaluronic acid inhibits interleukin-1-induced superoxide anion in bovine chondrocytes. *Inflamm Res* 1997 Mar; 46(3):114-117.
- 3.13 Ghatak S, Misra S, Toole BP. Hyaluronan oligosaccharides inhibit anchorage-independent growth of tumor cells by suppressing the phosphoinositide 3-kinase/Akt cell survival pathway. *J Biol Chem* 2002 Oct 11; 277(41):38013-38020.
- 3.14 Ghosh P. The role of hyaluronic acid (hyaluronan) in health and disease: interactions with cells, cartilage and components of synovial fluid. *Clin Exp Rheumatol* 1994 Jan-Feb; 12(1):75-82.
- 3.15 Ghosh P, Guidolin D. Potential mechanism of action of intra-articular hyaluronan therapy in osteoarthritis: are the effects molecular weight dependent? *Semin Arthritis Rheum* 2002 Aug; 32(1):10-37.
- 3.16 Goldberg RL, Toole BP. Hyaluronate inhibition of cell proliferation. *Arthritis Rheum* 1987 Jul; 30(7):769-778.
- 3.17 Gotoh S, Onaya J, Abe M, Miyazaki K, Hamai A, Horie K, Tokuyasu K. Effects of the molecular weight of hyaluronic acid and its action mechanisms on experimental joint pain in rats. *Ann Rheum Dis* 1993 Nov; 52(11):817-822.
- 3.18 Greco RM, Iacono JA, Ehrlich HP. Hyaluronic acid stimulates human fibroblast proliferation within a collagen matrix. *J Cell Physiol* 1998 Dec; 177(3):465-473.
- 3.19 Haslinger B, Mandl-Weber S, Sellmayer A, Sitter T. Hyaluronan fragments induce the synthesis of MCP-1 and IL-8 in cultured human peritoneal mesothelial cells. *Cell Tissue Res* 2001 Jul; 305(1):79-86.

- 3.20 Henderson EB, Grootveld M, Farrell A, Smith EC, Thompson PW, Blake DR. A pathological role for damaged hyaluronan in synovitis. *Ann Rheum Dis* 1991 Mar; 50(3):196-200.
- 3.21 Hodge-Dufour J, Noble PW, Horton MR, Bao C, Wysoka M, Burdick MD, Strieter RM, Trinchieri G, Pure E. Induction of IL-12 and chemokines by hyaluronan requires adhesion-dependent priming of resident but not elicited macrophages. *J Immunol* 1997 Sep 1; 159(5):2492-2500.
- 3.22 Homandberg GA, Hui F, Wen C, Kuettner KE, Williams JM. Hyaluronic acid suppresses fibronectin fragment mediated cartilage chondrolysis: I. In vitro *Osteoarthritis Cartilage* 1997 Sep; 5(5):309-319.
- 3.23 Horton MR, Shapiro S, Bao C, Lowenstein CJ, Noble PW. Induction and regulation of macrophage metalloelastase by hyaluronan fragments in mouse macrophages. *J Immunol* 1999 Apr 1; 162(7):4171-4176.
- 3.24 Huang L, Cheng YY, Koo PL, Lee KM, Qin L, Cheng JC, Kumta SM. The effect of hyaluronan on osteoblast proliferation and differentiation in rat calvarial-derived cell cultures. *J Biomed Mater Res A* 2003 Sep 15; 66(4):880-884.
- 3.25 Ialenti A, Di Rosa M. Hyaluronic acid modulates acute and chronic inflammation. *Agents Actions* 1994 Nov; 43(1-2):44-47.
- 3.26 Kang Y, Eger W, Koepf H, Williams JM, Kuettner KE, Homandberg GA. Hyaluronan suppresses fibronectin fragment-mediated damage to human cartilage explant cultures by enhancing proteoglycan synthesis. *J Orthop Res* 1999 Nov; 17(6):858-869.
- 3.27 Kawasaki K, Ochi M, Uchio Y, Adachi N, Matsusaki M. Hyaluronic acid enhances proliferation and chondroitin sulfate synthesis in cultured chondrocytes embedded in collagen gels. *J Cell Physiol* 1999 May; 179(2):142-148.
- 3.28 Knudson CB. Hyaluronan receptor-directed assembly of chondrocyte pericellular matrix. *J Cell Biol* 1993 Feb; 120(3):825-834.
- 3.29 Knudson CB, Knudson W. Hyaluronan and CD44: modulators of chondrocyte metabolism. *Clin Orthop* 2004 Oct; (427 Suppl):S152-S162.
- 3.30 Knudson W, Loeser RF. CD44 and integrin matrix receptors participate in cartilage homeostasis. *Cell Mol Life Sci* 2002 Jan; 59(1):36-44.
- 3.31 Knudson W, Casey B, Nishida Y, Eger W, Kuettner KE, Knudson CB. Hyaluronan oligosaccharides perturb cartilage matrix homeostasis and induce chondrocytic chondrolysis. *Arthritis Rheum* 2000 May; 43(5):1165-1174.
- 3.32 Kobayashi H, Suzuki M, Kanayama N, Nishida T, Takigawa M, Terao T. CD44 stimulation by fragmented hyaluronic acid induces upregulation of urokinase-type plasminogen activator and its receptor and subsequently facilitates invasion of human chondrosarcoma cells. *Int J Cancer* 2002 Dec 1; 102(4):379-389.
- 3.33 Kurosaka N, Takagi T, Koshino T. Effects of hyaluronate on CD44 expression of infiltrating cells in exudate of rat air pouch, induced by sensitization with lipopolysaccharide. *J Rheumatol* 1999 Oct; 26(10):2186-2190.
- 3.34 Larsen NE, Lombard KM, Parent EG, Balazs EA. Effect of hylan on cartilage and chondrocyte cultures. *J Orthop Res* 1992 Jan; 10(1):23-32.
- 3.35 Lees VC, Fan TP, West DC. Angiogenesis in a delayed revascularization model is accelerated by angiogenic oligosaccharides of hyaluronan. *Lab Invest* 1995 Aug; 73(2):259-266.
- 3.36 McKee CM, Penno MB, Cowman M, Burdick MD, Strieter RM, Bao C, Noble PW. Hyaluronan (HA) fragments induce chemokine gene expression in alveolar macrophages. The role of HA size and CD44. *J Clin Invest* 1996 Nov 15; 98(10):2403-2413.
- 3.37 McKee CM, Lowenstein CJ, Horton MR, Wu J, Bao C, Chin BY, Choi AM, Noble PW. Hyaluronan fragments induce nitric-oxide synthase in murine macrophages through a nuclear factor kappaB-dependent mechanism. *J Biol Chem* 1997 Mar 21; 272(12):8013-8018.
- 3.38 Moreland LW. Intra-articular hyaluronan (hyaluronic acid) and hylans for the treatment of osteoarthritis: mechanisms of action. *Arthritis Res Ther* 2003; 5(2):54-67.
- 3.39 Moseley R, Leaver M, Walker M, Waddington RJ, Parsons D, Chen WY, Embury G. Comparison of the antioxidant properties of HYAFF-11p75, AQUACEL and hyaluronan towards reactive oxygen species in vitro. *Bio-materials* 2002 May; 23(10):2255-2264.
- 3.40 Nakamura K, Yokohama S, Yoneda M, Okamoto S, Tamaki Y, Ito T, Okada M, Aso K, Makino I. High, but not low, molecular weight hyaluronan prevents T-cell-mediated liver injury by reducing proinflammatory cytokines in mice. *J Gastroenterol* 2004; 39(4):346-354.
- 3.41 Neumann A, Schnitzel R, Palm D, Riederer P, Munch G. High molecular weight hyaluronic acid inhibits advanced glycation endproduct-induced NF-kappaB activation and cytokine expression. *FEBS Lett* 1999 Jun 25; 453(3):283-287.

- 3.42 Noble PW. Hyaluronan and its catabolic products in tissue injury and repair. *Matrix Biol* 2002 Jan; 21(1):25-29.
- 3.43 Noble PW, Lake FR, Henson PM, Riches DW. Hyaluronate activation of CD44 induces insulin-like growth factor-1 expression by a tumor necrosis factor-alpha-dependent mechanism in murine macrophages. *J Clin Invest* 1993; 91:2368-2377.
- 3.44 Noble PW, McKee CM, Cowman M, Shin HS. Hyaluronan fragments activate an NF-kappa B/I-kappa B autoregulatory loop in murine macrophages. *J Exp Med* 1996; 183:2373-2378.
- 3.45 Ohkawara Y, Tamura G, Iwasaki T, Tanaka A, Kikuchi T, Shirato K. Activation and transforming growth factor-beta production in eosinophils by hyaluronan. *Am J Respir Cell Mol Biol* 2000 Oct; 23(4):444-451.
- 3.46 Ohno-Nakahara M, Honda K, Tanimoto K, Tanaka N, Doi T, Suzuki A, Yoneno K, Nakatani Y, Ueki M, Ohno S, Knudson W, Knudson CB, Tanne K. Induction of CD44 and MMP expression by hyaluronidase treatment of articular chondrocytes. *J Biochem (Tokyo)* 2004 May; 135(5):567-575.
- 3.47 Peluso GF, Perbellini A, Tajana GF. The effect of high and low molecular weight hyaluronic acid on mitogen-induced lymphocyte proliferation. *Curr Ther Res* 1990; 47:437-443.
- 3.48 Pisko EJ, Turner RA, Soderstrom LP, Panetti M, Foster SL, Treadway WJ. Inhibition of neutrophil phagocytosis and enzyme release by hyaluronic acid. *Clin Exp Rheumatol* 1983 Jan-Mar; 1(1):41-44.
- 3.49 Presti D, Scott JE. Hyaluronan-mediated protective effect against cell damage caused by enzymatically produced hydroxyl (OH) radicals is dependent on hyaluronan molecular mass. *Cell Biochem Funct* 1994 Dec; 12(4):281-288.
- 3.50 Punzi L. The complexity of the mechanisms of action of hyaluronan in joint diseases. *Clin Exp Rheumatol* 2001 May-Jun; 19(3):242-246.
- 3.51 Rockey CD, Chung JJ, McKee CM, Noble PW. Stimulation of inducible nitric oxide synthase in rat liver by hyaluronan fragments. *Hepatology* 1998 Jan; 27(1):86-92.
- 3.52 Rossler A, Hinghofer-Szalkay H. Hyaluronan fragments: an information-carrying system? *Horm Metab Res* 2003 Feb; 35(2):67-68.
- 3.53 Sattar A, Kumar S, West DC. Does hyaluronan have a role in endothelial cell proliferation of the synovium? *Semin Arthritis Rheum* 1992 Aug; 22(1):37-43.
- 3.54 Scott JE. Supramolecular organization of extracellular matrix glycosaminoglycans, in vitro and in the tissues. *FASEB J* 1992 Jun; 6(9):2639-2645.
- 3.55 Scott JE, Heatley F. Biological properties of hyaluronan in aqueous solution are controlled and sequestered by reversible tertiary structures, defined by NMR spectroscopy. *Biomacromolecules* 2002 May-Jun; 3(3):547-553.
- 3.56 Scott JE, Heatley F. Biological properties of hyaluronan are controlled and sequestered by tertiary structures. In *Hyaluronan Volume 1 Chemical, Biochemical and Biological Aspects* Kennedy JF, Phillips GO, Williams PA, Hascall VC, Eds., Woodhead Publishing Ltd., Cambridge, England; 2002:117-122.
- 3.57 Slevin M, Krupinski J, Kumar S, Gaffney J. Angiogenic oligosaccharides of hyaluronan induce protein tyrosine kinase activity in endothelial cells and activate a cytoplasmic signal transduction pathway resulting in proliferation. *Lab Invest* 1998 Aug; 78(8):987-1003.
- 3.58 Smith MM, Ghosh P. The synthesis of hyaluronic acid by human synovial fibroblasts is influenced by the nature of the hyaluronate in the extracellular environment. *Rheumatol Int* 1987; 7(3):113-122.
- 3.59 Stern R. Hyaluronan catabolism: a new metabolic pathway. *Eur J Cell Biol* 2004 Aug; 83(7):317-325.
- 3.60 Suzuki M, Kobayashi H, Kanayama N, Nishida T, Takigawa M, Terao T. CD44 stimulation by fragmented hyaluronic acid induces upregulation and tyrosine phosphorylation of c-Met receptor protein in human chondrosarcoma cells. *Biochim Biophys Acta* 2002 Aug 19; 1591(1-3):37-44.
- 3.61 Tanaka M, Masuko-Hongo K, Kato T, Nishioka K, Nakamura H. Suppressing effects of hyaluronan on MMP-1 and RANTES production from chondrocytes. *Rheumatol Int* 2004 Dec 3, [Epub ahead of print].
- 3.62 Termeer CC, Hennies J, Voith U, Ahrens T, Weiss JM, Prehm P, Simon JC. Oligosaccharides of hyaluronan are potent activators of dendritic cells. *J Immunol* 2000 Aug 15; 165(4):1863-1870.
- 3.63 Termeer C, Benedix F, Sleeman J, Fieber C, Voith U, Ahrens T, Miyake K, Freudenberg M, Galanos C, Simon JC. Oligosaccharides of hyaluronan activate dendritic cells via toll-like receptor 4. *J Exp Med* 2002 Jan 7; 195(1):99-111.
- 3.64 Termeer CC, Prehm P, Simon JC. Low molecular weight oligosaccharides of hyaluronan potentially activate dendritic cells. In *Hyaluronan. Clinical, Biochemical and Biological Aspects*, Vol 1, Kennedy JF, Phillips GO, Williams PA, Hascall VC, Eds., Woodhead Publishing Limited, Cambridge, UK, 2002, 457-468.
- 3.65 Tobetto K, Nakai K, Akatsuka M, Yasui T, Ando T, Hirano S. Inhibitory effects of hyaluronan on neutrophil-mediated cartilage degradation. *Connect Tissue Res* 1993; 29(3):181-190.

- 3.66 Tobetto K, Yasui T, Ando T, Hayaishi M, Motohashi N, Shinogi M, Mori I. Inhibitory effects of hyaluronan on [¹⁴C]arachidonic acid release from labeled human synovial fibroblasts. *Jpn J Pharmacol* 1992 Oct; 60(2):79-84.
- 3.67 Toole BP. Hyaluronan: from extracellular glue to pericellular cue. *Nat Rev Cancer* 2004 Jul; 4(7):528-539.
- 3.68 Walsh DA. Angiogenesis and arthritis. *Rheumatology* 1999 Feb; 38(2):103-112.
- 3.69 Weigel PH, Fuller GM, LeBoeuf RD. A model for the role of hyaluronic acid and fibrin in the early events during the inflammatory response and wound healing. *J Theor Biol* 1986 Mar 21; 119(2):219-234.
- 3.70 West DC, Kumar S. The effect of hyaluronate and its oligosaccharides on endothelial cell proliferation and monolayer integrity. *Exp Cell Res* 1989 Jul; 183(1):179-196.
- 3.71 West DC, Hampson IN, Arnold F, Kumar S. Angiogenesis induced by degradation products of hyaluronic acid. *Science* 1985 Jun 14; 228(4705):1324-1326.
- 3.72 Wiebkin OW, Hardingham TE, Muir H. The interaction of proteoglycans and hyaluronic acid and the effect of hyaluronic acid on proteoglycan synthesis by chondrocytes of adult cartilage. Ch 4 in *Dynamics of Connective Tissue Macromolecules*. Burleigh PMC, Poole AR, Eds., North-Holland Publishing Company, Amsterdam, 1975, 81-104.
- 3.73 Xu H, Ito T, Tawada A, Maeda H, Yamanokuchi H, Isahara K, Yoshida K, Uchiyama Y, Asari A. Effect of hyaluronan oligosaccharides on the expression of heat shock protein 72. In *Hyaluronan. Clinical, Biochemical and Biological Aspects*, Vol 1, Kennedy JF, Phillips GO, Willaims PA, Hascall VC, Eds., Woodhead Publishing Limited, Cambridge, UK, 2002, 435-442.
- 3.74 Yasui T, Akatsuka M, Tobetto K, Hayaishi M, Ando T. The effect of hyaluronan on interleukin-1 alpha-induced prostaglandin E2 production in human osteoarthritic synovial cells. *Agents Actions* 1992 Sep; 37(1-2):155-156.
- 3.75 Yasui T, Akatsuka M, Tobetto K, Umemoto J, Ando T, Yamashita K, Hayakawa T. Effects of hyaluronan on the production of stromelysin and tissue inhibitor of metalloproteinase-1 (TIMP-1) in bovine articular chondrocytes. *Biomed Res* 1992; 13(5):343-348.

control, toxicological testing and reproducibility than the other two types of HA material.

As mentioned previously in this review, HA with high molecular weight of 1 million daltons has viscosity and lubricating properties, and sends antiinflammatory, anabolic signals to cells. Indeed, one mechanism of action proposed for intraarticular HA treatments of OA is the normalization of cell signaling to the synovium from high molecular weight HA (66, 71). Table 3 lists some of the biological properties associated with high molecular weight HA as well as HA fragments. Products using high molecular weight HA match the identity of native HA in the body and from the scientific literature. Claims relating properties and actions of HA from products containing high molecular weight HA appropriately borrow evidence from the scientific literature. Thus, high molecular weight, purified, native HA is the most logical choice for use as HA material in dietary supplements.

ORAL ABSORPTION OF HA

Of course, one prerequisite for any beneficial effects from HA in dietary supplement products is uptake into bloodstream and joint tissues after oral intake. Of the three types of HA materials, only one type, purified HA, has published evidence of oral absorption. An abstract from the 2004 FASEB meeting in Washington DC described uptake of labeled HA into the bloodstream and localization of the label into the joints and salivary glands of rats and dogs several hours after oral administration (157). The HA in this study was derived from microbial fermentation, and was 1 million daltons. Since ^{99m}technetium was used as the label, and control experiments showed that the label was not removed or exchanged from HA molecules (Bucci, L., unpublished data), the presence of the label in the joints is most likely explained by the presence of the HA absorbed after oral

administration. This study is the first published report of oral absorption and tissue uptake of oral HA and provides the rationale for HA in dietary supplements.

Another source of purified HA from rooster comb has stated on the company website that absorption of their material has been found (158). However, no details or data have been presented or published. Hydrolyzed chicken sternal cartilage material is likewise claimed to have not only absorption, but superior absorption to native HA because of its smaller size. This claim is also on product labels or accompanying text (see table 2 and previous sections of this review). Again, website information from the manufacturer describes that absorption was found in a study, but no details or data have been presented or published in a scientific forum. No head-to-head comparisons to other HA materials concerning uptake have been reported, rendering claims of superior absorption unsubstantiated and unsupported. Only one type of HA—purified, high molecular weight HA from microbial fermentation—has published evidence for oral uptake and distribution to connective tissues.

BENEFICIAL PROPERTIES OF HA FOR JOINT HEALTH

Oral HA products are mostly targeted to joint and skin health, according to structure and function claims on product labels and accompanying literature (for a sampling, see Table 2). These two uses coincide with the prevalence of HA in the human body (8). Almost all beneficial properties for joint health attributable to HA have been documented for high molecular weight HA unbound to protein. Extensive *in vitro* studies using purified HA have uncovered beneficial properties for joints (see table 3). In addition, publication of a large number of human clinical studies using intraarticular injection of HA has demonstrated direct benefits for joint health (reviewed in 21, 22, 25, 26, 68-86).

Of particular interest are the human studies showing repair or normalization of synovium and cartilage structure and architecture for long time periods after a course of injections to persons with OA (66, 71, 159-164). These changes were documented by before and after tissue biopsies, along with biochemical measurements long after the injected HA disappeared from joints. These studies show direct evidence in humans that presence of purified, high molecular weight HA in joints has long-term benefits months after exogenous HA was given. The results of the oral uptake study presented at FASEB (157) show that oral HA has the potential to reach joints, and provide a strong rationale for use of oral HA in dietary supplements. However, at this time, this rationale applies only to purified, high molecular weight HA from microbial fermentation sources.

SUMMARY

It is interesting to speculate that continued oral administration of HA might produce similar or superior effects to intermittent injectable HA, as has been seen for both glucosamine and chondroitin sulfate. However, dietary supplements containing HA are not equivalent due to the inherent properties of the three major types of HA commercially available as dietary supplement materials. Consumers and health care professionals need to be aware of the different types of HA and their very large differences in properties (even before ingestion). One source, hydrolyzed chicken sternal cartilage, is clearly unlike native HA, does not match the biological properties of native HA, and consequently should not be represented as HA to consumers on product labels. Rooster comb HA is significantly smaller than native HA (about 10-20% the size of native HA), and is tightly bound to connective tissue proteins. This type of HA does not match the literature on properties and benefits of native HA because of its smaller size and tight binding to proteins. Finally, high molecu-

lar weight, purified HA is available from vegetarian-compatible and animal sources, virtually identical to native HA and pharmaceutical preparations of HA. This material is analogous to the HA used to generate the extensive body of knowledge on HA, both its roles and its therapeutic effects in OA. HA from the microbial fermentation method of preparation has published evidence for uptake into joints after oral administration. From all considerations, purified, high molecular weight HA from microbial fermentation in dietary supplement products has the highest likelihood for matching the known attributes of HA. The real HA—purified, high molecular weight—can stand up now.

REFERENCES

- Guss JM, Hukins DW, Smith PJ, Winter WT, Arnott S. Hyaluronic acid: molecular conformations and interactions in two sodium salts. *J Mol Biol* 1975 Jul 5; 95(3):359-384.
- Cleland RL, Sherblom AP. Isolation and physical characterization of hyaluronic acid prepared from bovine nasal septum by cetylpyridinium chloride precipitation. *J Biol Chem* 1977 Jan 25; 252(2):420-426.
- Shimada E, Matsumura G. Thin-layer chromatography of hyaluronate oligosaccharides. *J Biochem (Tokyo)* 1984 Sep; 96(3):721-725.
- Dahl LB, Dahl IM, Engstrom-Laurent A, Granath K. Concentration and molecular weight of sodium hyaluronate in synovial fluid in patients with rheumatoid arthritis and other arthropathies. *Ann Rheum Dis* 1985 Dec; 44(12):817-822.
- Hakansson L, Venge P. The molecular basis of the hyaluronic acid-mediated stimulation of granulocyte function. *J Immunol* 1987 Jun 15; 138(12):4347-4352.
- Laurent TC. Biochemistry of hyaluronan. *Acta Otolaryngol Suppl* 1987; 442:7-24.
- Holmes MW, Bayliss MT, Muir H. Hyaluronic acid in human articular cartilage. Age-related changes in content and size. *Biochem J* 1988 Mar; 250(2):435-441.
- Laurent TC, Fraser JR. Hyaluronan. *FASEB J* 1992 Apr; 6(7): 2397-2404.
- Kvam C, Granese D, Flaibani A, Zanetti F, Paoletti S. Purification and characterization of hyaluronan from synovial fluid. *Anal Biochem* 1993 May 15; 211(1):44-49.
- Lee HG, Cowman MK. An agarose gel electrophoretic method for analysis of hyaluronan molecular weight distribution. *Anal Biochem* 1994 Jun; 219(2):278-287.
- Ghosh P. The role of hyaluronic acid (hyaluronan) in health and disease: interactions with cells, cartilage and components of synovial fluid. *Clin Exp Rheumatol* 1994 Jan-Feb; 12(1):75-82.
- Laurent TC, Laurent UB, Fraser JR. Functions of hyaluronan. *Ann Rheum Dis* 1995 May; 54(5):429-432.
- Laurent TC, Laurent UB, Fraser JR. The structure and function of hyaluronan: An overview. *Immunol Cell Biol* 1996 Apr; 74(2):A1-A7.
- Fraser JRE, Laurent TC, Laurent UBG. Hyaluronan: its nature, distribution, functions and turnover. *J Intern Med* 1997 Jul; 242:27-33.
- Radin EL, Paul IL, Swann DA, Schottstaedt ES. Lubrication of synovial membrane. *Ann Rheum Dis* 1971 May; 30(3):322-325.
- Radin EL, Paul IL. A consolidated concept of joint lubrication. *J Bone Joint Surg* 1972; 54A:607-616.
- Swann DA, Radin EL, Nazimec M, Weisser PA, Curran N, Lewinnek G. Role of hyaluronic acid in joint lubrication. *Ann Rheum Dis* 1974; 33:318-326.
- Bowers WH. Lubrication of human joints. Ch 18 in *The Musculoskeletal System. Basic Processes and Disorders*, 2nd ed., Wilson FC, Ed., JB Lippincott Company, Philadelphia, 1983, 225-230.
- Varma R, Varma RS. Synovial fluid. Ch 12 in *Mucopolysaccharides - Glycosaminoglycans - of Body Fluids in Health and Disease*. Walter de Gruyter, Berlin, 1983, 345-370.
- Balazs EA, Denlinger JL. Clinical uses of hyaluronan. In: *The Biology of Hyaluronan*. John Wiley & Sons Ltd.: Sussex, UK, 1989; 265-275, discussion 275-280, 281-285.
- Balazs EA, Denlinger JL. Viscosupplementation: a new concept in the treatment of osteoarthritis. *J Rheumatol Suppl* 1993 Aug; 39:3-9.
- Pelletier JP, Martel-Pelletier J. The pathophysiology of osteoarthritis and the implication of the use of hyaluronan and hylan as therapeutic agents in viscosupplementation. *J Rheumatol Suppl* 1993 Aug; 39:19-24.
- McCarty DJ. Synovial fluid. Ch 5 in *Arthritis and Allied Conditions. A Textbook of Rheumatology*. Vol. 1, 12th ed., McCarty DJ, Koopman WJ, Eds., Lea & Febiger, Philadelphia, 1993, 63-84.

24. Simkin PA. Synovial physiology. Ch 10 in *Arthritis and Allied Conditions. A Textbook of Rheumatology*. Vol. 1, 12th ed., McCarty DJ, Koopman WJ, Eds., Lea & Febiger, Philadelphia, 1993, 199-212.
25. Aviad AD, Haupt JB. The molecular weight of therapeutic hyaluronan (sodium hyaluronate): how significant is it? *J Rheumatol* 1994 Feb; 21(2):297-301.
26. Goa KL, Benfield P. Hyaluronic acid. A review of its pharmacology and use as a surgical aid in ophthalmology, and its therapeutic potential in joint disease and wound healing. *Drugs* 1994 Mar; 47(3):536-566.
27. Sledge CB. Biology of the joint. Ch 1 in *Arthritis Surgery*: Sledge CB, Harris ED, Ruddy S, Kelley WN, Eds., WB Saunders, Philadelphia, 1994, 1-21.
28. Engström-Laurent A. Hyaluronan in joint disease. *J Intern Med* 1997 Jul; 242(1):57-60.
29. Lu Y, Levick JR, Wang W. Concentration polarisation of hyaluronan on the surface of the synovial lining of infused joints. *J Physiol* 2004 Oct 7 [Epub]
30. Rosenberg L. Structure of cartilage proteoglycans. Ch 5 in *Dynamics of Connective Tissue Macromolecules*. Burleigh PMC, Poole AR, Eds., North-Holland Publishing Company, Amsterdam, 1975, 105-128.
31. Muir IHM. Biochemistry. Ch 3 in *Adult Articular Cartilage*, 2nd ed., Freeman MAR, Ed., Pitman Medical Publishing Co. Ltd., Kent, UK, 1979, 145-214.
32. Roden L. Structure and metabolism of connective tissue proteoglycans. Ch 7 in *The Biochemistry of Glycoproteins and Proteoglycans*, Lennarz WJ, Ed., Plenum Press, New York, 1980, 267-372.
33. Hascall VC, Hascall GK. Proteoglycans. Ch 2 in *Cell Biology of Extracellular Matrix*, Hay ED, Ed., Plenum Press, New York, 1981, 39-64.
34. Bayliss MT. Proteoglycans: structure and molecular organisation in cartilage. Ch 3 in *Connective Tissue Matrix*. Hukins DWL, Ed., Macmillan, London, 1984, 55-88.
35. Caplan AI. Cartilage. *Sci Am* 1984 Oct; 251(4):84-97.
36. Heinegard D, Paulsson M. Structure and metabolism of proteoglycans. Ch 8 in *Extracellular Matrix Biochemistry*. Piez KA, Reddi AH, Eds., Elsevier, New York, 1984, 277-328.
37. Hardingham TE. Structure and biosynthesis of proteoglycans. Ch 6 in *Connective Tissue: Biological and Clinical Aspects*. Kuhn K, Krieg T, Eds., Karger, Basel, 1986, 143-183.
38. Prehm P. Mechanism, localization, and inhibition of hyaluronate synthesis. In *Articular Cartilage Biochemistry*. Kuettner KE, Schleyerbach R, Hascall VC, Eds., Raven Press, New York, 1986, 81-92.
39. Hunziker EB, Schenk RK. Structural organization of proteoglycans in cartilage. In *Biology of Proteoglycans*. Wight TN, Mecham RP, Eds., Academic Press, 1987, 155-186.
40. Silver FH. Self-assembly of connective tissue macromolecules. Ch 5 in *Biological Materials: Structure, Mechanical Properties, and Modeling of Soft Tissues*, New York University Press, New York, 1987, 136-163.
41. Gamble JG. Structural components. Ch 4 in *The Musculoskeletal System. Physiological Basics*. Raven Press, New York, 1988, 57-80.
42. Handley CJ, Ng CK. Proteoglycan, hyaluronan, and noncollagenous matrix protein metabolism by chondrocytes. Ch 3 in *Biological Regulation of the Chondrocytes*, Adolphe M, Ed., CRC Press, Boca Raton, 1992, 85-104.
43. Perkins SJ, Nealis AS, Dunham DG, Hardingham TE, Muir IH. Neutron and X-ray solution-scattering studies of the ternary complex between proteoglycan-binding region, link protein and hyaluronan. *Biochem J* 1992 Jul 1; 285(Pt 1):263-268.
44. Poole AR. Cartilage in health and disease. Ch 15 in *Arthritis and Allied Conditions. A Textbook of Rheumatology*. Vol. 1, 12th ed., McCarty DJ, Koopman WJ, Eds., Lea & Febiger, Philadelphia, 1993, 279-333.
45. Rosenberg L. Structure and function of proteoglycans. Ch 12 in *Arthritis and Allied Conditions. A Textbook of Rheumatology*. Vol. 1, 12th ed., McCarty DJ, Koopman WJ, Eds., Lea & Febiger, Philadelphia, 1993, 229-242.
46. Morgelin M, Heingard D, Engel J, Paulsson M. The cartilage proteoglycan aggregate: assembly through combined protein-carbohydrate and protein-protein interactions. *Biophys Chem* 1994 May; 50(1-2):113-128.
47. Melrose J, Little CB, Ghosh P. Detection of aggregatable proteoglycan populations by affinity blotting using biotinylated hyaluronan. *Anal Biochem* 1998 Feb 15; 256(2):149-157.
48. Hardingham TE. Cartilage: aggrecan-link protein-hyaluronan aggregates. www.glycoforum.gr.jp/science/hyaluronan/HA05/HA05E.html, March 15, 2002.
49. Yasumoto T, Bird JL, Sugimoto K, Mason RM, Bayliss MT. The G1 domain of aggrecan released from porcine articular cartilage forms stable com-

- plexes with hyaluronan/link protein. *Rheumatology (Oxford)* 2003 Feb; 42(2):336-342.
50. Green SJ, Tarone G, Underhill CB. Distribution of hyaluronate and hyaluronate receptors in the adult lung. *J Cell Sci* 1988 May; 90(Pt 1):145-156.
51. Raja RH, McGary CT, Weigel PH. Affinity and distribution of surface and intracellular hyaluronic acid receptors in isolated rat liver endothelial cells. *J Biol Chem* 1988 Nov 15; 263(32):16661-16668.
52. Alho AM, Underhill CB. The hyaluronate receptor is preferentially expressed on proliferating epithelial cells. *J Cell Biol* 1989 Apr; 108(4):1557-1565.
53. Wang C, Tammi M, Tammi R. Distribution of hyaluronan and its CD44 receptor in the epithelia of human skin appendages. *Histochemistry* 1992 Sep; 98(2):105-112.
54. Wheatley SC, Isacke CM, Crossley PH. Restricted expression of the hyaluronan receptor, CD44, during postimplantation mouse embryogenesis suggests key roles in tissue formation and patterning. *Development* 1993 Oct; 119(2):295-306.
55. Kennel SJ, Lankford TK, Foote LJ, Shipcock SG, Stringer C. CD44 expression on murine tissues. *J Cell Sci* 1993 Feb; 104(Pt 2):373-382.
56. Ranganathan S, Ganguly AK, Datta K. Evidence for presence of hyaluronan binding protein on spermatozoa and its possible involvement in sperm function. *Mol Reprod Dev* 1994 May; 38(1):69-76.
57. Behzad F, Seif MW, Campbell S, Aplin JD. Expression of two isoforms of CD44 in human endometrium. *Biol Reprod* 1994 Oct; 51(4):739-747.
58. Yasaka N, Furue M, Tamaki K. CD44 expression in normal human skin and skin tumors. *J Dermatol* 1995 Feb; 22(2):88-94.
59. Entwistle J, Hall CL, Turley EA. HA receptors: regulators of signaling to the cytoskeleton. *J Cell Biochem* 1996 Jun 15; 61(4):569-577.
60. Ponta H, Wainwright D, Herrlich P. The CD44 protein family. *Int J Biochem Cell Biol* 1998 Mar; 30(3):299-305.
61. Goodison S, Urquidí V, Tarin D. CD44 cell adhesion molecules. *Mol Pathol* 1999 Aug; 52(4):189-196.
62. Punzi L. The complexity of the mechanisms of action of hyaluronan in joint diseases. *Clin Exp Rheumatol* 2001 May-Jun; 19(3):242-246.
63. Ghosh P, Guidolin D. Potential mechanism of action of intra-articular hyaluronan therapy in osteoarthritis: are the effects molecular weight dependent? *Semin Arthritis Rheum* 2002 Aug; 32(1):10-37.
64. Turley EA, Noble PW, Bourguignon LY. Signaling properties of hyaluronan receptors. *J Biol Chem* 2002 Feb 15; 277(7):4589-4592.
65. Rossler A, Hinghofer-Szalkay H. Hyaluronan fragments: an information-carrying system? *Horm Metab Res* 2003 Feb; 35(2):67-68.
66. Uebelhart D, Williams JM. Effects of hyaluronic acid on cartilage degradation. *Curr Opin Rheumatol* 1999 Sep; 11(5):427-435.
67. Moreland LW. Intra-articular hyaluronan (hyaluronic acid) and hylans for the treatment of osteoarthritis: mechanisms of action. *Arthritis Res Ther* 2003; 5(2):54-67.
68. Simon LS. Viscosupplementation therapy with intra-articular hyaluronic acid. Fact or fantasy? *Rheum Dis Clin North Am* 1999 May; 25(2):345-357.
69. Adams ME, Lussier AJ, Peyron JG. A risk-benefit assessment of injections of hyaluronan and its derivatives in the treatment of osteoarthritis of the knee. *Drug Saf* 2000 Aug; 23(2):115-130.
70. Brandt KD, Smith GN Jr, Simon LS. Intraarticular injection of hyaluronan as treatment for knee osteoarthritis: what is the evidence? *Arthritis Rheum* 2000 Jun; 43(6):1192-1203.
71. Dougados M. Sodium hyaluronate therapy in osteoarthritis: arguments for a potential beneficial structural effect. *Semin Arthritis Rheum* 2000 Oct; 30(2 Suppl 1):19-25.
72. Hochberg MC. Role of intra-articular hyaluronic acid preparations in medical management of osteoarthritis of the knee. *Semin Arthritis Rheum* 2000 Oct; 30(2 Suppl 1):2-10.
73. Marshall KW. Intra-articular hyaluronan therapy. *Curr Opin Rheumatol* 2000 Sep; 12(5):468-474.
74. Moskowitz RW. Hyaluronic acid supplementation. *Curr Rheumatol Rep* 2000 Dec; 2(6):466-471.
75. Wen DY. Intra-articular hyaluronic acid injections for knee osteoarthritis. *Am Fam Physic* 2000 Aug 1; 62:565-570, 572.
76. Wright KE, Maurer SG, Di Cesare PE. Viscosupplementation for osteoarthritis. *Am J Orthop* 2000 Feb; 29(2):80-88.
77. Ayral X. Injections in the treatment of osteoarthritis. *Best Pract Res Clin Rheumatol* 2001 Oct; 15(4):609-626.
78. Kirwan J. Is there a place for intra-articular hyaluronate in osteoarthritis of the knee? *Knee* 2001 Jun; 8(2):93-101.
79. Pandolfi S, Exer P, Schwarz HA. Viscosupplementa-

- tion in arthrosis. *Ther Umsch* 2002 Oct; 59(10):545-549.
80. Altman RD. Status of hyaluronan supplementation therapy in osteoarthritis. *Curr Rheumatol Rep* 2003 Feb; 5(1):7-14.
81. Kelly MA, Goldberg VM, Healy WL, Pagnano MW, Hamburger MI. Osteoarthritis and beyond: a consensus on the past, present, and future of hyaluronans in orthopedics. *Orthopedics* 2003 Oct; 26(10):1064-1079; quiz 1080-1081.
82. Lo GH, LaValley M, McAlindon T, Felson DT. Intra-articular hyaluronic acid in treatment of knee osteoarthritis: a meta-analysis. *JAMA* 2003 Dec 17; 290(23):3115-3121.
83. Aggarwal A, Sempowski IP. Hyaluronic acid injections for knee osteoarthritis. Systematic review of the literature. *Can Fam Physician* 2004 Feb; 50:249-256.
84. Kelly MA, Kurzweil PR, Moskowitz RW. Intra-articular hyaluronans in knee osteoarthritis: rationale and practical considerations. *Am J Orthop* 2004 Feb; 33(2 Suppl):15-22.
85. Kelly MA, Moskowitz RW, Lieberman JR. Hyaluronan therapy: looking toward the future. *Am J Orthop* 2004 Feb; 33(2 Suppl):23-28.
86. Wang CT, Lin J, Chang CJ, Lin YT, Hou SM. Therapeutic effects of hyaluronic acid on osteoarthritis of the knee. A meta-analysis of randomized controlled trials. *J Bone Joint Surg Am* 2004 Mar; 86-A(3):538-545.
87. Kawcak CE, Frisbie DD, Trotter GW, McIlwraith CW, Gillette SM, Powers BE, Walton RM. Effects of intravenous administration of sodium hyaluronate on carpal joints in exercising horses after arthroscopic surgery and osteochondral fragmentation. *Am J Vet Res* 1997 Oct; 58(10):1132-1140.
88. Campo GM, Avenoso A, Campo S, Ferlazzo A, Altavilla D, Micali C, Calatroni A. Aromatic trap analysis of free radicals production in experimental collagen-induced arthritis in the rat: protective effect of glycosaminoglycans treatment. *Free Radic Res* 2003 Mar; 37(3):257-268.
89. Chapman C, Ehrhardt K, Blumenstein R. Efficacy of treatment with sodium hyaluronate on joints in collagen-induced arthritis in DBA/1J mice. *FASEB J* 2004 Mar; 15:A32. [abstract #68.4]
90. Pierce SW. Efficacy of orally administered sodium hyaluronate gel in the racing thoroughbred. In Ch 6, Musculoskeletal system, Hyaluronan 2003 Proceedings, Cleveland, OH, October 11-16, 2003. (www.matrixbiologyinstitute.org/ha03/)
91. Stancikova M, Svik K, Istok R, Rovensky J, Velebny V. The effects of hyaluronan on bone resorption and bone mineral density in a rat model of estrogen deficiency-induced osteopenia. *Int J Tiss react* 2004; 26(1-2):9-16.
92. Efficacy data, Contipro Group Holding, http://www.cpn-contipro.com/cpn/index.php?action=content&link=cpn_nutr&lang=cz.
93. Pavelka K, Manopulo R, Bucsi L. Double-blind, dose-effect study of oral Chondroitin 4&6 Sulfate 1200 mg, 800 mg, 200 mg and placebo in the treatment of knee osteoarthritis. *Litera Rheumatologica* 1999; 24:21-30.
94. Lee JY, Spicer AP. Hyaluronan: a multifunctional, megaDalton, stealth molecule. *Curr Opin Cell Biol* 2000 Oct; 12(5):581-586.
95. Camenisch TD, McDonald JA. Hyaluronan: is bigger better? *Am J Respir Cell Mol Biol* 2000 Oct; 23(4):431-433.
96. Noble PW. Hyaluronan and its catabolic products in tissue injury and repair. *Matrix Biol* 2002 Jan; 21(1):25-29.
97. Knudson CB, Knudson W. Hyaluronan and CD44: modulators of chondrocyte metabolism. *Clin Orthop* 2004 Oct; (427 Suppl):S152-S162.
98. Maneiro E, de Andres MC, Fernandez-Sueiro JL, Galdo F, Blanco FJ. The biological action of hyaluronan on human osteoarthritic articular chondrocytes: the importance of molecular weight. *Clin Exp Rheumatol* 2004 May-Jun; 22(3):307-312.
99. Ropes MM, Bauer W. Synovial fluid changes and joint disease. Cambridge, MA: Harvard University Press; 1953
100. Balazs EA, Watson D, Duff IF, Roseman S. Hyaluronic acid in synovial fluid. I. Molecular parameters of hyaluronic acid in normal and arthritic human fluids. *Arthritis Rheum* 1967 Aug; 10(4):357-376.
101. Bothner H, Wik O. Rheology of hyaluronate. *Acta Otolaryngol* (Stockh) 1987; Suppl. 442:25-30.
102. Jay GD, Haberstroh K, Cha CJ. Comparison of the boundary-lubricating ability of bovine synovial fluid, lubricin and Healon. *J Biomed Mater Res* 1998 Jun 5; 40(3):414-418.
103. Wobig M, Bach G, Beks P, Dickhut A, Runzheimer J, Schwieger G, Vetter G, Balazs E. The role of elastoviscosity in the efficacy of viscosupplementation for osteoarthritis of the knee: a comparison of hylan G-F 20 and lower-molecular-weight hyaluronan. *Clin Ther* 1999 Sep; 21(9):1549-1562.

104. Mabuchi K, Obara T, Ikegami K, Yamaguchi T, Kanayama T. Molecular weight independence of the effect of additive hyaluronic acid on the lubricating characteristics in synovial joints with experimental deterioration. *Clin Biomech (Bristol, Avon)* 1999 Jun; 14(5):352-356.
105. Mori S, Naito M, Moriyama S. Highly viscous sodium hyaluronate and joint lubrication. *Int Orthop* 2002; 26(2):116-121.
106. Mazzucco D, Scott R, Spector M. Composition of joint fluid in patients undergoing total knee replacement and revision arthroplasty: correlation with flow properties. *Biomaterials* 2004; 25:4433-4445.
107. Underhill CB. The interaction of hyaluronate with the cell surface: the hyaluronate receptor and the core protein. *Ciba Found Symp* 1989; 143:87-99; discussion 100-106, 281-285.
108. Aruffo A, Stamenkovic I, Melnick M, Underhill CB, Seed B. CD44 is the principal cell surface receptor for hyaluronate. *Cell* 1990 Jun 29; 61(7):1303-1313.
109. Fujii K, Tanaka Y, Hubscher S, Saito K, Ota T, Eto S. Crosslinking of CD44 on rheumatoid synovial cells augment interleukin 6 production. *Lab Invest* 1999 Dec; 79(12):1439-1446.
110. Ohkawara Y, Tamura G, Iwasaki T, Tanaka A, Kikuchi T, Shirato K. Activation and transforming growth factor-beta production in eosinophils by hyaluronan. *Am J Respir Cell Mol Biol* 2000 Oct; 23(4):444-451.
111. Haslinger B, Mandl-Weber S, Sellmayer A, Sitter T. Hyaluronan fragments induce the synthesis of MCP-1 and IL-8 in cultured human peritoneal mesothelial cells. *Cell Tissue Res* 2001 Jul; 305(1):79-86.
112. Kobayashi H, Suzuki M, Kanayama N, Nishida T, Takigawa M, Terao T. CD44 stimulation by fragmented hyaluronic acid induces upregulation of urokinase-type plasminogen activator and its receptor and subsequently facilitates invasion of human chondrosarcoma cells. *Int J Cancer* 2002 Dec 1; 102(4):379-389.
113. Huang L, Cheng YY, Koo PL, Lee KM, Qin L, Cheng JC, Kumta SM. The effect of hyaluronan on osteoblast proliferation and differentiation in rat calvarial-derived cell cultures. *J Biomed Mater Res A* 2003 Sep 15; 66(4):880-884.
114. Wang MJ, Jeng KC, Kuo JS, Chen HL, Huang HY, Chen WF, Lin SZ. C-Jun N-terminal kinase and, to a lesser extent, p38 mitogen-activated protein kinase regulate inducible nitric oxide synthase expression in hyaluronan fragments-stimulated BV-2 microglia. *J Neuroimmunol* 2004 Jan; 146(1-2):50-62.
115. Ito T, Williams JD, Al-Assaf S, Phillips GO, Phillips AO. Hyaluronan and proximal tubular cell migration. *Kidney Int* 2004 Mar; 65(3):823-833.
116. Nakamura K, Yokohama S, Yoneda M, Okamoto S, Tamaki Y, Ito T, Okada M, Aso K, Makino I. High, but not low, molecular weight hyaluronan prevents T-cell-mediated liver injury by reducing proinflammatory cytokines in mice. *J Gastroenterol* 2004; 39(4):346-354.
117. Toole BP. Hyaluronan: from extracellular glue to pericellular cue. *Nat Rev Cancer* 2004 Jul; 4(7):528-539.
118. Gotoh S, Onaya J, Abe M, Miyazaki K, Hamai A, Horie K, Tokuyasu K. Effects of the molecular weight of hyaluronic acid and its action mechanisms on experimental joint pain in rats. *Ann Rheum Dis* 1993 Nov; 52(11):817-822.
119. Asari A, Miyauchi S, Matsuzaka S, Ito T, Komiyama E, Uchiyama Y. Molecular weight-dependent effects of hyaluronate on the arthritic synovium. *Arch Histol Cytol* 1998 May; 61(2):125-135.
120. Tikiz C, Unlu Z, Sener A, Efe M, Tuzun C. Comparison of the efficacy of lower and higher molecular weight viscosupplementation in the treatment of hip osteoarthritis. *Clin Rheumatol* 2005 Jan 13; [Epub ahead of print].
121. Hodge-Dufour J, Noble PW, Horton MR, Bao C, Wysoka M, Burdick MD, Strieter RM, Trinchieri G, Pure E. Induction of IL-12 and chemokines by hyaluronan requires adhesion-dependent priming of resident but not elicited macrophages. *J Immunol* 1997 Sep 1; 159(5):2492-2500.
122. Alaniz L, Cabrera PV, Blanco G, Ernst G, Rimoldi G, Alvarez E, Hajos SE. Interaction of CD44 with different forms of hyaluronic acid. Its role in adhesion and migration of tumor cells. *Cell Commun Adhes* 2002 May-Jun; 9(3):117-130.
123. Ghatak S, Misra S, Toole BP. Hyaluronan oligosaccharides inhibit anchorage-independent growth of tumor cells by suppressing the phosphoinositide 3-kinase/Akt cell survival pathway. *J Biol Chem* 2002 Oct 11; 277(41):38013-38020.
124. Toole BP, Wight TN, Tammi MI. Hyaluronan-cell interactions in cancer and vascular disease. *J Biol Chem* 2002 Feb; 277(7):4593-4596.
125. Fieber C, Baumann P, Vallon R, Termeer C, Simon

- JC, Hofmann M, Angel P, Herrlich P, Sleeman JP. Hyaluronan-oligosaccharide-induced transcription of metalloproteases. *J Cell Sci* 2004 Jan 15; 117(Pt 2):359-367.
126. Kinoshita M, Kakehi K. Analysis of the interaction between hyaluronan and hyaluronan-binding proteins by capillary affinity electrophoresis: significance of hyaluronan molecular size on binding reaction. *J Chromatogr B Analyt Technol Biomed Life Sci* 2005 Feb 25; 816(1-2):289-295.
 127. Rooney P, Kumar S, Ponting J, Wang M. The role of hyaluronan in tumour neovascularization (review). *Int J Cancer* 1995 Mar 3; 60(5):632-636.
 128. Scott JE, Heatley F. Biological properties of hyaluronan are controlled and sequestered by tertiary structures. In *Hyaluronan Volume 1 Chemical, Biochemical and Biological Aspects* Kennedy JF, Phillips GO, Williams PA, Hascall VC, Eds., Woodhead Publishing Ltd., Cambridge, England; 2002:117-122.
 129. Suzuki M, Kobayashi H, Kanayama N, Nishida T, Takigawa M, Terao T. CD44 stimulation by fragmented hyaluronic acid induces upregulation and tyrosine phosphorylation of c-Met receptor protein in human chondrosarcoma cells. *Biochim Biophys Acta* 2002 Aug 19; 1591(1-3):37-44.
 130. Sugahara KN, Murai T, Hirata T, Miyasaka M. Hyaluronan oligosaccharides induce CD44 cleavage and promote cell migration in CD44-expressing tumor cells. In Ch. 4, Musculoskeletal System, *Hyaluronan 2003 Proceedings*, Cleveland, OH, October 11-16, 2003. (www.matrixbiologyinstitute.org/ha03)
 131. Sugahara KN, Murai T, Nishinakamura H, Kawashima H, Saya H, Miyasaka M. Hyaluronan oligosaccharides induce CD44 cleavage and promote cell migration in CD44-expressing tumor cells. *J Biol Chem* 2003 Aug 22; 278(34):32259-32265.
 132. Stern R. Hyaluronan catabolism: a new metabolic pathway. *Eur J Cell Biol* 2004 Aug; 83(7):317-325.
 133. Forrester JV, Balazs EA. Inhibition of phagocytosis by high molecular weight hyaluronate. *Immunology* 1980 Jul; 40(3):435-436.
 134. West DC, Kumar S. The effect of hyaluronate and its oligosaccharides on endothelial cell proliferation and monolayer integrity. *Exp Cell Res* 1989 Jul; 183(1):179-196.
 135. McKee CM, Penno MB, Cowman M, Burdick MD, Strieter RM, Bao C, Noble PW. Hyaluronan (HA) fragments induce chemokine gene expression in alveolar macrophages. The role of HA size and CD44. *J Clin Invest* 1996 Nov 15; 98(10):2403-2413.
 136. McKee CM, Lowenstein CJ, Horton MR, Wu J, Bao C, Chin BY, Choi AM, Noble PW. Hyaluronan fragments induce nitric-oxide synthase in murine macrophages through a nuclear factor kappaB-dependent mechanism. *J Biol Chem* 1997 Mar 21; 272(12):8013-8018.
 137. Rockey CD, Chung JJ, McKee CM, Noble PW. Stimulation of inducible nitric oxide synthase in rat liver by hyaluronan fragments. *Hepatology* 1998 Jan; 27(1):86-92.
 138. Horton MR, Shapiro S, Bao C, Lowenstein CJ, Noble PW. Induction and regulation of macrophage metalloelastase by hyaluronan fragments in mouse macrophages. *J Immunol* 1999 Apr 1; 162(7):4171-4176.
 139. Blass SL, Pure E, Hunter CA. A role for CD44 in the production of IFN-gamma and immunopathology during infection with *Toxoplasma gondii*. *J Immunol* 2001 May 1; 166(9):5726-5732.
 140. Ohno-Nakahara M, Honda K, Tanimoto K, Tanaka N, Doi T, Suzuki A, Yoneno K, Nakatani Y, Ueki M, Ohno S, Knudson W, Knudson CB, Tanne K. Induction of CD44 and MMP expression by hyaluronidase treatment of articular chondrocytes. *J Biochem (Tokyo)* 2004 May; 135(5):567-575.
 141. Tanaka M, Masuko-Hongo K, Kato T, Nishioka K, Nakamura H. Suppressing effects of hyaluronan on MMP-1 and RANTES production from chondrocytes. *Rheumatol Int* 2004 Dec 3, [Epub ahead of print].
 142. Noble PW, Lake FR, Henson PM, Riches DW. Hyaluronate activation of CD44 induces insulin-like growth factor-1 expression by a tumor necrosis factor-alpha-dependent mechanism in murine macrophages. *J Clin Invest* 1993; 91:2368-2377.
 143. Noble PW, McKee CM, Cowman M, Shin HS. Hyaluronan fragments activate an NF-kappa B/I-kappa B autoregulatory loop in murine macrophages. *J Exp Med* 1996; 183:2373-2378.
 144. Henderson EB, Grootveld M, Farrell A, Smith EC, Thompson PW, Blake DR. A pathological role for damaged hyaluronan in synovitis. *Ann Rheum Dis* 1991 Mar; 50(3):196-200.
 145. Ishaq S. Hyaluronic acid and chondroitin sulfate based hydrolyzed collagen type II and method of making same. United States Patent 6,780,841, August 24, 2004.
 146. Alkalayi A. Hydrolyzed collagen type II and uses

- thereof. United States Patent 6, 025,327, February 15, 2000.
147. Alkalayi A. Method of making hydrolyzed collagen type II. United States Patent 6, 323, 319, November 27, 2001.
 148. Kalman DS, Schwartz HI, Pachon J, Sheldon E, Almada AL. A randomized double blind clinical pilot trial evaluating the safety and efficacy of hydrolyzed collagen type II in adults with osteoarthritis. *FASEB J* 2004 Apr; 18(4):A90 [abstract LB435].
 149. Biocell Technology. Clinical study shows Biocell Collagen II effective. Press Release, April 5, 2003 from Biocell website. (www.biocelltechnology.com/press5.htm)
 150. Biocell Technology. Clinical study shows hyaluronic acid in Biocell Collagen II® found to have significant absorption and bioavailability. Press Release, February 2, 2004 from Biocell website (www.biocelltechnology.com/press9.htm).
 151. Rossler A, Laszlo Z, Kvas E, Hinghofer-Szalkay HG. Plasma hyaluronan concentration: no circadian rhythm but large effect of food intake in humans. *Eur J Appl Physiol Occup Physiol* 1998 Nov; 78(6):573-577.
 152. Udell RG, Naguib YMA. Hyaluronic acid in soft gel form. United States Patent 6,806,259, October 19, 2004.
 153. Swann DA. Studies on hyaluronic acid. II. The protein component(s) of rooster comb hyaluronic acid. *Biochim Biophys Acta* 1968 May 6; 160(1):96-105
 154. Swann DA, Caulfield JB. Studies on hyaluronic acid. V. Relationship between the protein content and viscosity of rooster comb dermis hyaluronic acid. *Connect Tissue Res* 1975; 4(1):31-39.
 155. Tsiganos CP, Vynios DH, Kalpaxis DL. Rooster comb hyaluronate-protein, a non-covalently linked complex. *Biochem J* 1986 Apr 1; 235(1):117-123.
 156. Hyalgan® Prescribing Information. http://www.sanori-synthelabo.us/products/pi_hyalgan/pi_hyalgan.html
 157. Schauss AG, Balogh L, Polyak A, Mathe D, Kiraly R, Janoki G. Absorption, distribution and excretion of ^{99m}technetium labeled hyaluronan after single oral doses in rats and beagle dogs. *FASEB J* 2004 Apr; 18(4), A150-A151. [abstract 129.4]
 158. Bioiberica website for Hyal-Joint: <http://www.bioiberica.com/jointcare/hyaljoint.htm>
 159. Dougados M, Ayrat X, Listrat V, Bonvarlet JP, Simonnet J, Amor B. La chondroscopie: un nouvel instrument de mesure de l'arthrose? [Chondroscopy: a new method for measuring osteoarthritis?] *Rev Rhum Ed Fr* 1994 Nov 15; 61(9 Pt 2):131S-136S.
 160. Listrat V, Ayrat X, Patarnello F, Bonvarlet JP, Simonnet J, Amor B, Dougados M. Arthroscopic evaluation of potential structure modifying activity of hyaluronan (Hyalgan) in osteoarthritis of the knee. *Osteoarthritis Cartilage* 1997 May; 5(3):153-160.
 161. Frizziero L, Govoni E, Bacchini P. Intra-articular hyaluronic acid in the treatment of osteoarthritis of the knee: clinical and morphological study. *Clin Exp Rheumatol* 1998 Jul-Aug; 16(4):441-449.
 162. Pasquali Ronchetti I, Guerra D, Taparelli F, Boraldi F, Bergamini G, Mori G, Zizzi F, Frizziero L. Morphological analysis of knee synovial membrane biopsies from a randomized controlled clinical study comparing the effects of sodium hyaluronate (Hyalgan) and methylprednisolone acetate (Depomedrol) in osteoarthritis. *Rheumatology (Oxford)* 2001 Feb; 40(2):158-169.
 163. Guidolin DD, Ronchetti IP, Lini E, Guerra D, Frizziero L. Morphological analysis of articular cartilage biopsies from a randomized, clinical study comparing the effects of 500-730 kDa sodium hyaluronate (Hyalgan) and methylprednisolone acetate on primary osteoarthritis of the knee. *Osteoarthritis Cartilage* 2001 May; 9(4):371-381.
 164. Jubb RW, Piva S, Beinat L, Dacre J, Gishen P. A one-year, randomised, placebo (saline) controlled clinical trial of 500-730 kDa sodium hyaluronate (Hyalgan) on the radiological change in osteoarthritis of the knee. *Int J Clin Pract* 2003 Jul-Aug; 57(6):467-474.